

Spectral decomposition of some non-self-adjoint operators

Jérémy Faupin

Université de Lorraine, Metz

Mathematical aspects of the physics with non-self-adjoint operators
CIRM, June 2024

J.F. and **N. Frantz**, *Spectral decomposition of some non-self-adjoint operators*, Annales Henri Lebesgue, 6, 1115-1167, (2023).

Abstract model

The model

- \mathcal{H} complex Hilbert space
- Hamiltonian

$$H = H_0 + V = H_0 + CWC,$$

with $H_0 \geq 0$, $C \in \mathcal{B}(\mathcal{H})$, $C > 0$ and relatively compact with respect to H_0 , $W \in \mathcal{B}(\mathcal{H})$ arbitrary

- H is a **closed** operator with domain

$$\mathcal{D}(H) = \mathcal{D}(H_0)$$

- $-iH$ generates a strongly continuous group $\{e^{-itH}\}_{t \in \mathbb{R}}$ s.t.

$$\|e^{-itH}\| \leq e^{\|V\||t|}, \quad t \in \mathbb{R}$$

- $H^* = H_0 + CW^*C$ with domain $\mathcal{D}(H^*) = \mathcal{D}(H_0)$
- $\sigma_{\text{ess}}(H) = \sigma_{\text{ess}}(H_0)$ and $\sigma(H) \setminus \sigma_{\text{ess}}(H)$ consists of an **at most countable number of eigenvalues of finite algebraic multiplicities** that can only accumulate at points of $\sigma_{\text{ess}}(H)$




FIGURE: Spectrum of H

Point spectral subspace (I)

Point spectrum

$$\sigma_p(H) := \{\lambda \in \mathbb{C}, \text{Ker}(H - \lambda) \neq \{0\}\}$$

Algebraic multiplicity of an eigenvalue $\lambda \in \sigma_p(H)$:

$$m_\lambda := \dim\left(\bigcup_{k \geq 1} \text{Ker}((H - \lambda)^k)\right)$$

Discrete spectrum and discrete spectral subspace

$$\sigma_{\text{disc}}(H) := \sigma(H) \setminus \sigma_{\text{ess}}(H) \subset \sigma_p(H)$$

- For $\lambda \in \sigma_{\text{disc}}(H)$, **Riesz projection** defined by

$$\Pi_\lambda = -\frac{1}{2i\pi} \int_\gamma R_H(z) dz, \quad R_H(z) = (H - z)^{-1},$$

where γ is a circle centered at λ , of sufficiently small radius

- $\text{Ran}(\Pi_\lambda)$ spanned by generalized eigenvectors of H associated to λ , $u \in \mathcal{D}(H^k)$ s.t. $(H - \lambda)^k u = 0$
- Discrete spectral subspace:

$$\mathcal{H}_{\text{disc}}(H) := \text{Span} \{u \in \text{Ran}(\Pi_\lambda), \lambda \in \sigma_{\text{disc}}(H)\}^{\text{cl}}$$

Point spectral subspace (II)

Set of embedded eigenvalues

$$\sigma_{\text{emb}}(H) := \sigma_{\text{p}}(H) \cap \sigma_{\text{ess}}(H)$$

To define spectral projections corresponding to embedded eigenvalues

- We suppose the existence of an anti-linear continuous map $J : \mathcal{H} \rightarrow \mathcal{H}$ satisfying

$$J\mathcal{D}(H_0) \subset \mathcal{D}(H_0) \quad \text{and} \quad \forall u \in \mathcal{D}(H_0), \quad JHu = H^*Ju$$

- If $\lambda \in \sigma_{\text{emb}}(H)$, we suppose that $m_\lambda < \infty$ and that the symmetric bilinear form

$$\text{Ker}((H - \lambda)^{m_\lambda}) \ni (u, v) \mapsto \langle Ju, v \rangle \quad \text{is non-degenerate}$$

- Under these conditions, there exists a basis $(\varphi_k)_{1 \leq k \leq m_\lambda}$ of $\text{Ker}((H - \lambda)^{m_\lambda})$ such that $\langle J\varphi_i, \varphi_j \rangle = \delta_{ij}$, $1 \leq i, j \leq m_\lambda$. Then

$$\Pi_\lambda u := \sum_{k=1}^{m_\lambda} \langle J\varphi_k, u \rangle \varphi_k, \quad u \in \mathcal{H}$$

-

$$\mathcal{H}_{\text{emb}}(H) := \text{Span} \{u \in \text{Ran}(\Pi_\lambda), \lambda \in \sigma_{\text{emb}}(H)\}^{\text{cl}}$$

Point spectral subspace (III)

Point spectral subspace

$$\mathcal{H}_p(H) := \mathcal{H}_{\text{disc}}(H) \oplus \mathcal{H}_{\text{emb}}(H)$$

Asymptotically disappearing states


Subspaces of asymptotically disappearing states

$$\mathcal{H}_{\text{ads}}^{\pm}(H) := \left\{ u \in \mathcal{H}, \lim_{t \rightarrow \pm\infty} \|e^{-itH} u\| = 0 \right\}^{\text{cl}}$$

Relation with discrete generalized eigenstates

- Easy to see that

$$\text{Span} \{ \text{Gen. eigenstates associated to } \lambda, \pm \text{Im}(\lambda) < 0 \}^{\text{cl}} \subset \mathcal{H}_{\text{ads}}^{\pm}(H)$$



- **Question:** conditions implying that the previous inclusion becomes an equality?
- [Kato '66] For small perturbations, H and H_0 are similar, hence $\mathcal{H}_{\text{ads}}^{\pm}(H) = \{0\}$
- For dissipative operators, $\text{Im}(V) \leq 0$, the question was left as an open problem in [Davies '80], with an answer given in [F., Fröhlich '18]

Absolutely continuous spectral subspace

Absolutely continuous spectral subspace

$$\mathcal{H}_{\text{ac}}(H) := \left\{ u \in \mathcal{H}, \exists c_u > 0, \forall v \in \mathcal{H}, \int_{\mathbb{R}} |\langle e^{-itH} u, v \rangle|^2 dt \leq c_u \|v\|^2 \right\}^{\text{cl}}$$

Relation with point spectral subspace of H^*

- Not difficult to verify that

$$\mathcal{H}_{\text{ac}}(H) \subset \mathcal{H}_{\text{p}}(H^*)^\perp$$

- **Question:** conditions implying that the previous inclusion becomes an equality?
- Other definitions considered in the literature: [Davies '79] for dissipative operators, [Naboko '76] using the theory of dilations of dissipative operators
- Under suitable assumption, coincides with the space of 'scattering states'

Spectral singularities (I)


Definition: Spectral singularity

- 1 $\lambda \in \sigma_{\text{ess}}(H)$ is an **outgoing/incoming regular spectral point** of H if λ is not an accumulation point of eigenvalues located in $\lambda \pm i(0, \infty)$ and if the limit

$$CR_H(\lambda \pm i0^+)CW := \lim_{\varepsilon \rightarrow 0^+} CR_H(\lambda \pm i\varepsilon)CW$$

exists in the norm topology of $\mathcal{B}(\mathcal{H})$

- 2 λ is a regular spectral point of H if it is both an incoming and an outgoing regular spectral point of H
- 3 **Spectral singularity = not regular spectral point**



Spectral singularities (II)

Remarks

For dissipative operators, similar definition in [F, Fröhlich '18]. Other related notions:

- [Dunford '58] (theory of spectral operators), [Schwartz '59] (spectral singularity = singular point of a 'spectral resolution' for non-self-adjoint operators)
- [F, Nicoleau '19] (for dissipative operators, spectral singularity = point of the essential spectrum where the scattering matrix is non-invertible)
- For one-dimensional Schrödinger operators (spectral singularity = zero of the Jost function)

Some properties [F, Frantz]

- λ **embedded eigenvalue** $\Rightarrow \lambda$ both **outgoing and incoming spectral singularity**
- At **thresholds**, outgoing and incoming spectral singularities **coincide**

Spectral singularities (III)

Proposition (Schrödinger operators in 3-dimension)

Suppose that V is a complex-valued potential such that $\langle x \rangle^\sigma V(x) \in L^\infty(\mathbb{R}^3)$ with $\sigma > 1$. Let $C(x) = \langle x \rangle^{-\sigma/2}$. Then for all $\lambda > 0$, the following conditions are equivalent:

- 1 λ is an outgoing/incoming spectral singularity of $H = -\Delta + V$
- 2 There exists $\Psi \neq 0$, $\langle x \rangle^{-\sigma/2} \Psi \in L^2$, Ψ satisfying the outgoing/incoming Sommerfeld radiation condition, such that

$$(-\Delta + V(x) - \lambda)\Psi = 0$$

The same holds at the threshold $\lambda = 0$ if $\langle x \rangle^\sigma V(x) \in L^\infty(\mathbb{R}^3)$ with $\sigma > 2$

Remarks

- There is an abstract version of this proposition involving the **Gelfand triple**

$$\text{Ran}(C) \hookrightarrow \mathcal{H} \hookrightarrow (\text{Ran}(C))'$$

- [Wang '12]: For any $\lambda > 0$, one can construct a smooth compactly supported potential V such that λ is a **spectral singularity of H**

Hypotheses (I)

(H1) Limiting absorption principle for H_0

$$\sup_{\pm \text{Im}(z) > 0} \|CR_0(z)C\| < \infty$$

Remark

Satisfied for $H_0 = -\Delta$, $C(x) = \langle x \rangle^{-\sigma/2}$, $\sigma > 2$ in dimension 3

Consequences

- The spectrum of H_0 is **purely absolutely continuous**, i.e.

$$\sigma_{\text{pp}}(H_0) = \emptyset, \quad \sigma_{\text{ac}}(H_0) = \sigma(H_0), \quad \sigma_{\text{sc}}(H_0) = \emptyset$$

- The limits $CR_0(\lambda \pm i0^+)C$ exist for almost every $\lambda \in \sigma_{\text{ess}}(H)$, in the norm topology of $\mathcal{B}(\mathcal{H})$

Hypotheses (II)

(H2) Eigenvalues of H

H only has a **finite number of eigenvalues** with **finite algebraic multiplicities**

Remark

- Satisfied for Schrödinger operators $H = -\Delta + V(x)$ in $L^2(\mathbb{R}^3)$ if V is **exponentially decaying** [Frank, Laptev, Safronov '16]
- Does **not** exclude embedded eigenvalues


Hypotheses (III)

(H3) Spectral singularities for H

H only has a **finite number of spectral singularities** $\{\lambda_1, \dots, \lambda_n\} \subset \sigma_{\text{ess}}(H)$ and there exist $\varepsilon_0 > 0$, integers $\nu_1, \dots, \nu_n, -\nu_\infty \geq 0$ and neighborhoods $\mathcal{V}_1, \dots, \mathcal{V}_n, \mathcal{V}_\infty$ such that

$$\sup_{\text{Re}(z) \in \mathcal{V}_j, \pm \text{Im}(z) \in (0, \varepsilon_0)} |z - \lambda_j|^{\nu_j} \|CR_H(z)CW\| < \infty,$$

(with the convention $\lambda_\infty = 0$)



Remark

For Schrödinger operators $H = -\Delta + V(x)$ in $L^2(\mathbb{R}^3)$ with V **compactly supported**, (H3) is satisfied with ν_j the multiplicity of the resonance λ_j and $\nu_\infty = 0$

Asymptotically disappearing states

Recall that

$$\mathcal{H}_{\text{ads}}^{\pm}(H) := \left\{ u \in \mathcal{H}, \lim_{t \rightarrow \pm\infty} \|e^{-itH}u\| = 0 \right\}^{\text{cl}}$$

Theorem [F, Frantz]

Suppose that Hypotheses (H1)–(H3) hold. Then

$$\mathcal{H}_{\text{ads}}^{\pm}(H) = \text{Span} \{ \text{Gen. eigenstates associated to } \lambda, \pm \text{Im}(\lambda) < 0 \}^{\text{cl}}$$

Remark

For dissipative operators, analogous result proven in [F. Fröhlich '18]. The proof in [F. Fröhlich '18] relies on the existence and properties of wave operators. Our proof does not rely on scattering theory

Theorem (Consequence for Schrödinger operators)

Suppose that V is a complex-valued potential such that $V \in L_c^{\infty}(\mathbb{R}^3)$. Then the previous theorem applies to $H = -\Delta + V$.

Hypotheses (IV)

(H4) Embedded eigenvalues

There exists an **anti-linear continuous** map $J : \mathcal{H} \rightarrow \mathcal{H}$ such that

- 1 $J\mathcal{D}(H_0) \subset \mathcal{D}(H_0)$ and $\forall u \in \mathcal{D}(H_0), JH_0u = H_0Ju$
- 2 $JC = CJ$ and $JW = W^*J$

Moreover, for all embedded eigenvalues $\lambda \in \sigma_{\text{ess}}(H)$, the symmetric bilinear form

$$\text{Ker}((H - \lambda)^{m_\lambda}) \ni (u, v) \mapsto \langle Ju, v \rangle \text{ is non-degenerate}$$

Remark

For Schrödinger operators $H = -\Delta + V(x)$, J is the **complex conjugation** and Hypothesis (H4) means that for all embedded eigenvalues $\lambda \in [0, \infty)$,

$$\text{Ker}((H - \lambda)^{m_\lambda}) \ni (u, v) \mapsto \int_{\mathbb{R}^3} u(x)v(x)dx \text{ is non-degenerate}$$

Absolutely continuous spectral subspace

Recall that

$$\mathcal{H}_{\text{ac}}(H) := \left\{ u \in \mathcal{H}, \exists c_u > 0, \forall v \in \mathcal{H}, \int_{\mathbb{R}} |\langle e^{-itH} u, v \rangle|^2 dt \leq c_u \|v\|^2 \right\}^{\text{cl}}$$

Theorem [F, Frantz]

Suppose that Hypotheses (H1)–(H3) hold. If H has embedded eigenvalues, suppose in addition that (H4) holds. Then

$$\mathcal{H}_{\text{ac}}(H) = \mathcal{H}_{\text{p}}(H^*)^{\perp}$$

Remark: comparable results in the literature (only for H dissipative)

[Simon '79] for dissipative Schrödinger operators, [Davies '80] for abstract dissipative operators using the theory of dilations, with a different definition of \mathcal{H}_{ac} and a different result (\mathcal{H}_{ac} coincides with the orthogonal complement of 'bound states')

Theorem (Consequence for Schrödinger operators)

Suppose that V is a complex-valued potential such that $V \in L_c^\infty(\mathbb{R}^3)$ and the previous hypothesis on embedded eigenvalues is satisfied. Then the previous theorem applies to $H = -\Delta + V$

Spectral decomposition

Consequence of the previous two theorems

Suppose that Hypotheses (H1)–(H4) hold. Then we have the following ***J*-orthogonal direct sum** decompositions of the Hilbert space:

$$\begin{aligned}\mathcal{H} &= \mathcal{H}_{\text{ac}}(H) \oplus \mathcal{H}_{\text{p}}(H) \\ &= \mathcal{H}_{\text{ac}}(H) \oplus \mathcal{H}_{\text{disc}}(H) \oplus \mathcal{H}_{\text{emb}}(H) \\ &= \mathcal{H}_{\text{ac}}(H) \oplus \mathcal{H}_{\text{ads}}^+(H) \oplus \mathcal{H}_{\text{ads}}^-(H) \oplus \mathcal{H}_{\text{b}}(H),\end{aligned}$$

where $\mathcal{H}_{\text{b}}(H)$ is the space of '**bound states**', i.e. the closure of the vector space spanned by all generalized eigenvectors of H corresponding to real eigenvalues (either isolated or embedded)

Spectral
decomposition
of
some non-
self-adjoint
operators

Jérémy
Faupin

The
abstract
framework

Spectral de-
composition

Regularized
functional
calculus

Thank you !

Spectral
decomposition
of
some non-
self-adjoint
operators

Jérémy
Faupin

The
abstract
framework

Spectral de-
composition

Regularized
functional
calculus

Regularized functional calculus

Regularized Stone formula

Stone's formula for self-adjoint operators

Suppose H is a **self-adjoint** operator without embedded eigenvalues. Then

$$\text{Id} = \sum_{\lambda \in \sigma_{\text{disc}}(H)} \Pi_{\lambda} + \text{w-lim}_{\varepsilon \rightarrow 0^+} \frac{1}{2\pi i} \int_{\sigma_{\text{ess}}(H)} (R_H(\lambda + i\varepsilon) - R_H(\lambda - i\varepsilon)) d\lambda$$

Regularized version

Recall that we have assumed

$$\sup_{\substack{\text{Re}(z) \in \sigma_{\text{ess}}(H) \\ \pm \text{Im}(z) \in (0, \varepsilon_0)}} |r(z)| \|CR_H(z)CW\| < \infty, \quad r(z) = \frac{1}{(z - z_0)^{\nu_\infty}} \prod_{j=1}^n \frac{(z - \lambda_j)^{\nu_j}}{(z - z_0)^{\nu_j}}$$

Then, under our assumptions, we have

$$r(H) = \sum_{\lambda \in \sigma_{\text{disc}}(H)} r(H) \Pi_{\lambda} + \text{w-lim}_{\varepsilon \rightarrow 0^+} \frac{1}{2\pi i} \int_{\sigma_{\text{ess}}(H)} r(\lambda) (R_H(\lambda + i\varepsilon) - R_H(\lambda - i\varepsilon)) d\lambda$$

Idea of the proof

Riesz-Dunford functional calculus

$$r(H) = -\frac{1}{2i\pi} \int_{\Gamma_\varepsilon} r(z)R_H(z)dz \quad \text{then } \varepsilon \rightarrow 0^+$$




FIGURE: The contour Γ_ε .

Functional calculus (I)

Recall Hypothesis (H1):

$$\sup_{\pm \text{Im}(z) > 0} \|CR_0(z)C\| < \infty$$

Proposition (Functional calculus in intervals not containing spectral singularities)

Suppose (H1). Let $I \subset \mathbb{R}$ be a closed interval and suppose that there exists $\varepsilon_0 > 0$ such that

$$\sup_{\text{Re}(z) \in I, \pm \text{Im}(z) \in (0, \varepsilon_0)} \|CR_H(z)CW\| < \infty.$$

Then the map

$$C_b(I) \ni f \mapsto f(H) := \text{w-lim}_{\varepsilon \rightarrow 0^+} \frac{1}{2\pi i} \int_I f(\lambda) (R_H(\lambda + i\varepsilon) - R_H(\lambda - i\varepsilon)) d\lambda \in \mathcal{B}(\mathcal{H})$$

Remark

Related to the Dunford-Schwartz theory of spectral operators

Functional calculus (II)

Proposition (Regularized functional calculus)

Suppose (H1). Let $I \subset \mathbb{R}$ be a closed interval and suppose that there exists $\varepsilon_0 > 0$ and a **bounded holomorphic function** h such that

$$\sup_{\substack{\operatorname{Re}(z) \in I \\ \pm \operatorname{Im}(z) \in (0, \varepsilon_0)}} |h(z)| \|CR_H(z)CW\| < \infty, \quad \lambda \mapsto \sup_{0 < \varepsilon < \varepsilon_0} |h'(\lambda \pm i\varepsilon)| \in L^2(I).$$

Then the map

$$\begin{aligned} C_{b, \text{reg}}(I) \ni f \mapsto f(H) := & \text{w-lim}_{\varepsilon \rightarrow 0^+} \frac{1}{2\pi i} \int_I g(\lambda) (h(\lambda + i\varepsilon)R_H(\lambda + i\varepsilon) \\ & - h(\lambda - i\varepsilon)R_H(\lambda - i\varepsilon)) d\lambda \in \mathcal{B}(\mathcal{H}) \end{aligned}$$

is an algebra morphism and there exists $c > 0$ such that

$$\|f(H)\|_{\mathcal{B}(\mathcal{H})} \leq c \|g\|_{L^\infty}.$$

Here

$$C_{b, \text{reg}}(I) := \{f : I \rightarrow \mathbb{C}, \exists g \in C_b(I), f = hg\}$$

Remark

Other functional calculi for operators on Banach spaces under an assumption of **polynomial growth of the resolvent** near the real axis: [Davies '95] (general theory), [Georgescu, Gérard, Häfner '13] (Krein spaces)

Spectral singularities and resonances (I)

$\mathcal{H} = L^2(\mathbb{R}^d)$, $H_0 = -\Delta$, V compactly supported

Resonance may be defined as a **pole** of the map

$$\mathbb{C} \ni z \mapsto (H - z^2)^{-1} : L_c^2(\mathbb{R}^3) \rightarrow L_{\text{loc}}^2(\mathbb{R}^3),$$

Then

Spectral singularity at $\lambda > 0$ = resonance at $\pm \lambda^{1/2}$

Remarks

- Resonances theory: [Sjöstrand '02], [Dyatlov-Zworski '18]
- [Wang '12]: For any $\lambda > 0$, one can construct a smooth compactly supported potential V such that λ is a **spectral singularity of H**

Spectral singularities and resonances (II)

Example: $\mathcal{H} = L^2(\mathbb{R}^3)$, $H_0 = -\Delta$, V short-range

If $V(x) = \mathcal{O}(\langle x \rangle^{-\delta})$ with $\delta > 1$, then $\pm\lambda^{1/2}$ (with $\lambda > 0$) may be called a **resonance** of H if the equation $(H - \lambda)u = 0$ admits a distributional solution (called a **resonant state**)

$$u \in \bigcap_{\sigma > 1} L^2_{-\sigma/2} \setminus L^2$$

satisfying the **Sommerfeld radiation condition**

$$u(x) = |x|^{-1} e^{\pm i\lambda^{1/2}|x|} \left(a\left(\frac{x}{|x|}\right) + o(1) \right), \quad |x| \rightarrow \infty,$$

with $a \in L^2(S^2)$, $a \neq 0$. Here $L^2_{-\sigma/2} = \{f : \mathbb{R}^3 \rightarrow \mathbb{C}, x \mapsto \langle x \rangle^{-\frac{\sigma}{2}} f(x) \in L^2(\mathbb{R}^3)\}$

Spectral singularities: Characterization (I)

Assumption (given $\lambda \in \sigma_{\text{ess}}(H)$, with $R_0 := R_{H_0}$)

$$CR_0(\lambda \pm i0^+)C := \lim_{\varepsilon \rightarrow 0^+} CR_0(\lambda \pm i\varepsilon)C \text{ exist in the topology of } \mathcal{B}(\mathcal{H}) \quad (*)$$

Free Schrödinger operator in $\mathcal{H} = L^2(\mathbb{R}^3)$

- For $\lambda > 0$, the limits

$$\langle x \rangle^{-s} (-\Delta - (\lambda \pm i0^+)) \langle x \rangle^{-s},$$

exist in the norm topology of $\mathcal{B}(\mathcal{H})$, for any $s > \frac{1}{2}$, where $\langle x \rangle := (1 + x^2)^{\frac{1}{2}}$

- If $\lambda = 0$, the limits

$$\langle x \rangle^{-s} (-\Delta \pm i0^+) \langle x \rangle^{-s},$$

exist (and coincide) for any $s > 1$

Spectral singularities: Characterization (II)

Extension of the Hilbert space

- Let $\mathcal{H}_C := \text{Ran}(C)$ equipped with $\langle u, v \rangle_{\mathcal{H}_C} := \langle C^{-1}u, C^{-1}v \rangle$
- Let \mathcal{H}'_C be the anti-dual of \mathcal{H}_C . We obtain the Gelfand triple

$$\mathcal{H}_C \hookrightarrow \mathcal{H} \hookrightarrow \mathcal{H}'_C$$

- Assuming that $\mathcal{D}(H_0|_{\mathcal{H}_C}) := \{u \in \mathcal{D}(H_0) \cap \mathcal{H}_C, H_0u \in \mathcal{H}_C\}$ is dense in \mathcal{H}_C , H extends to

$$H' = H'_0 + CWC' : \mathcal{H}'_C \rightarrow \mathcal{H}'_C$$

- $CR_0(\lambda \pm i0^+)C := \lim_{\varepsilon \rightarrow 0^+} CR_0(\lambda \pm i\varepsilon)C$ exists in $\mathcal{B}(\mathcal{H})$ is equivalent to

$$R_0(\lambda \pm i0^+) := \lim_{\varepsilon \rightarrow 0^+} R_0(\lambda \pm i\varepsilon) \text{ exist in } \mathcal{B}(\mathcal{H}_C, \mathcal{H}'_C)$$

Incoming/outgoing resonant states

Let $\lambda \in \sigma_{\text{ess}}(H)$ be a spectral singularity of H . The space $\mathcal{H}_C^{\pm}(\lambda) \subset \mathcal{H}'_C$ of **outgoing/incoming resonant states** corresponding to λ is defined by

$$\mathcal{H}_C^{\pm}(\lambda) := \text{Ker}(\text{Id} + R_0(\lambda \pm i0^+)CWC')$$

Spectral singularities: Characterization (III)

Theorem [F, Frantz]

Suppose that $(*)$ holds. The following conditions are equivalent:

- 1 λ is an **outgoing/incoming spectral singularity of H**
- 2 λ is an **eigenvalue of H'** associated to an **eigenvector $\Psi \in \mathcal{H}'_C^\pm(\lambda)$**

Some consequences

- λ **embedded eigenvalue** $\Rightarrow \lambda$ both **outgoing and incoming spectral singularity**
- At **thresholds**, outgoing and incoming spectral singularities **coincide**
- Suppose that V is a complex-valued potential such that $\langle x \rangle^\sigma V(x) \in L^\infty(\mathbb{R}^3)$ with $\sigma > 1$. Let $C(x) = \langle x \rangle^{-\sigma/2}$. Then for all $\lambda > 0$, the following conditions are equivalent:

- 1 λ is an outgoing/incoming spectral singularity of H
- 2 There exists $\Psi \in \mathcal{H}'_C^\pm(\lambda) \subset L^2_{-\sigma/2}$, $\Psi \neq 0$, such that

$$(-\Delta + V(x) - \lambda)\Psi = 0$$

- The same holds at the threshold $\lambda = 0$ if $\langle x \rangle^\sigma V(x) \in L^\infty(\mathbb{R}^3)$ with $\sigma > 2$

Spectral singularities: Ingredient of the proof

Proposition (Birman-Schwinger principle for spectral singularities)

Suppose that $(*)$ holds. Then the following conditions are equivalent:

- 1 λ is an outgoing/incoming regular spectral point of H
- 2 $\text{Id} + CR_0(\lambda \pm i0^+)CW$ is invertible in $\mathcal{B}(\mathcal{H})$

Remark

Birman-Schwinger principle recently studied in abstract non-self-adjoint settings:

- [Behrndt, ter Elst, Gesztesy '20]
- [Hansmann, Krejcirik '20]